

# Solution to an Open Problem - A Characterization of Conditionally Cancellative T-subnorms

Balasubramaniam Jayaram

**Abstract.** In this work we solve an open problem of U.Höhle ([10], Problem 11). We show that the solution gives a characterization of all conditionally cancellative t-subnorms. Further, we give an equivalence condition for a conditionally cancellative t-subnorm to be a t-norm and hence show that conditionally cancellative t-subnorms whose natural negations are strong are, in fact, t-norms.

**Mathematics Subject Classification (2010).** 26D07, 39B62 (54E70).

**Keywords.** Triangular norms, conditional cancellativity, t-subnorm, fuzzy negation, natural negation, residuation, open problem.

## 1. Introduction

*”Triangular norms are, on the one hand, special semigroups and, on the other hand, solutions of some functional equations. This mixture quite often requires new approaches to answer questions about the nature of triangular norms.”* With this observation, Klement et al., in [10], present a collection of open problems posed during the 24<sup>th</sup> Linz Seminar on fuzzy set theory. They deal with unsolved problems (as of then) related to fuzzy aggregation operations, especially t-norms and t-subnorms. For other collections of open problems related to triangular norms and related operators, see [11, 12]

Solutions to these open problems involving triangular norms are of both theoretical and applicational interest. Recently solutions to quite a few open problems in the field of triangular norms have been published [2, 4, 8, 14, 17, 18, 21], including in this journal, see for instance, [7, 13, 20, 19]. Since the publication of [10], some problems mentioned therein have been solved - for instance, Problem 1 was solved by Ouyang et al. [16], Problem 5 was solved by Ouyang and Li [15] while for some other problems partial solutions have been given, see for instance, the papers of Viceník [22], [23], [24] relating to Problem 4(i).

One of the open problems listed therein was posed by Prof. U. Höhle (Problem 11) which reads as follows:

**Problem 1.1 (U.Höhle, [10], Problem 11).** *Characterize all left-continuous  $t$ -norms  $T$  which satisfy*

$$I(x, T(x, y)) = \max(n(x), y), \quad x, y \in [0, 1]. \quad (1.1)$$

where  $I$  is the residual operator linked to  $T$ , i.e.,

$$I(x, y) = \sup\{t \in [0, 1] | T(x, t) \leq y\}, \quad x, y \in [0, 1], \quad (1.2)$$

$$n(x) = n_T(x) = I(x, 0) \text{ for all } x \in [0, 1]. \quad (1.3)$$

Further, Prof. U.Höhle goes on to remark the following:

*Remark 1.2.* "In the class of continuous  $t$ -norms, only nilpotent  $t$ -norms fulfill the above property."

In this work we deal with two problems. Firstly, we solve the above open problem of U.Höhle and show that the solution gives a characterization of all conditionally cancellative  $t$ -subnorms. From the proven result it does follow that the remark of Prof. U.Höhle - Remark 1.2 - is not always true and give an equivalence condition for it to be true, viz., that the natural negation obtained from the  $t$ -norm is strong.

Secondly, this quite naturally leads us to consider conditionally cancellative  $t$ -subnorms whose natural negations are involutive. Once again, by proving an equivalence condition for a conditionally cancellative  $t$ -subnorm to be a  $t$ -norm, we show that conditionally cancellative  $t$ -subnorms whose natural negations are involutive, in fact, become  $t$ -norms.

## 2. Preliminaries

**Definition 2.1.** *A function  $N: [0, 1] \rightarrow [0, 1]$  is called a fuzzy negation if  $N$  is decreasing and  $N(0) = 1$ ,  $N(1) = 0$ .  $N$  is said to be involutive or strong if  $N \circ N = id_{[0,1]}$ .*

**Definition 2.2 ([9], Definition 1.7).** *A  $t$ -subnorm is a function  $M: [0, 1]^2 \rightarrow [0, 1]$  such that it is monotonic non-decreasing, associative, commutative and  $M(x, y) \leq \min(x, y)$  for all  $x, y \in [0, 1]$ .*

Note that for a  $t$ -subnorm 1 need not be the neutral element, unlike in the case of a  $t$ -norm.

**Definition 2.3 (cf. [9], Definition 2.9 (iii)).** *A  $t$ -subnorm  $M$  satisfies the Conditional Cancellation Law if, for any  $x, y, z \in (0, 1)$ ,*

$$M(x, y) = M(x, z) > 0 \text{ implies } y = z. \quad (\text{CCL})$$

Alternately, (CCL) implies that on the positive domain of  $M$ , i.e., on the set  $\{(x, y) \in (0, 1]^2 \mid M(x, y) > 0\}$ ,  $M$  is strictly increasing. See Figure 1 (a) and (b) for examples of a conditionally cancellative  $t$ -subnorm and one that is not.




FIGURE 1. While  $M_{\mathbf{Pf}}$  is a conditionally cancellative t-subnorm,  $M_{\mathbf{B}}$  is not.

**Definition 2.4** (cf. [1], **Definition 2.3.1**). *Let  $M$  be any t-subnorm. Its natural negation  $n_M$  is given by*

$$n_M(x) = \sup\{t \in [0, 1] \mid M(x, t) = 0\}, \quad x \in [0, 1]. \quad (2.1)$$

Note that though  $n_M(0) = 1$ , it need not be a fuzzy negation, since  $n_M(1)$  can be greater than 0. However, we have the following result.

**Lemma 2.5** (cf. [1], **Proposition 2.3.4**). *Let  $M$  be any t-subnorm and  $n_M$  its natural negation. Then we have the following:*

- (i)  $M(x, y) = 0 \implies y \leq n_M(x)$ .
- (ii)  $y < n_M(x) \implies M(x, y) = 0$ .
- (iii) *If  $M$  is left-continuous then  $y = n_M(x) \implies M(x, y) = 0$ , i.e., the reverse implication of (i) also holds.*

*Proof.* Let  $\mathcal{A}_x = \{t \in [0, 1] \mid M(x, t) = 0\}$ . Clearly,  $0 \in \mathcal{A}_x$  since  $M(x, 0) = 0$ . Moreover, by the monotonicity of  $M$  we have that if  $t \in \mathcal{A}_x$  then  $[0, t] \subseteq \mathcal{A}_x$ . Thus, for every  $x \in (0, 1)$  we have that  $\mathcal{A}_x = [0, \alpha_x)$  or  $[0, \alpha_x]$  for some  $\alpha_x \in [0, 1]$ . Now,  $n_M(x) = \sup \mathcal{A}_x = \alpha_x$ .

- (i) Let  $M(x, y) = 0$ . Then  $y \in \mathcal{A}_x$  and hence  $y \leq \sup \mathcal{A}_x = n_M(x)$ .
- (ii) Conversely, let  $y < \sup \mathcal{A}_x = n_M(x)$ . Then  $y \in \mathcal{A}_x$  and hence  $M(x, y) = 0$ .
- (iii) If  $M$  is left-continuous then  $y = n_M(x) = \alpha_x \in \mathcal{A}_x$  and  $M(x, y) = 0$ . The converse follows from (i).

□

### 3. Solution to the Open Problem of U. Höhle

It should be noted that in the case  $T$  is left-continuous - as stated in **Problem 1** - the sup in (1.2) actually becomes max. It is worth mentioning that the residual can be determined for more generalised conjunctions and the conditions underwhich this residual becomes a fuzzy implication can be found in, for instance, [3, 5]. Hence we further generalise the statement of **Problem 1** by considering a t-subnorm instead of a t-norm and also dropping the condition of left-continuity. As we show below the solution characterizes the set of all conditionally cancellative t-subnorms.

**Theorem 3.1.** *Let  $M$  be any t-subnorm and  $I$  the residual operation linked to  $M$  by (1.2). Then the following are equivalent:*

- (i) *The pair  $(I, M)$  satisfies (1.1).*
- (ii)  *$M$  is a Conditionally Cancellative t-subnorm.*

*Proof.* Let  $M$  be any t-subnorm, not necessarily left-continuous. Note that we denote  $n_M$  simply by  $n$ .

- (i)  $\implies$  (ii): Let the adjoint pair  $(I, M)$  satisfy (1.1). On the contrary, let us assume that there exist  $x, y, z \in (0, 1)$  such that  $M(x, y) = M(x, z) > 0$  but  $y < z$ . Then we have that

$$\text{LHS (1.1)} = I(x, M(x, y)) = \sup\{t \in [0, 1] \mid M(x, t) \leq M(x, y)\} \geq z > y .$$

However, note that, from Lemma 2.5 (i) we have that  $y \geq n(x)$ , since  $M(x, y) > 0$ . Thus

$$\text{RHS (1.1)} = \max(n(x), y) = y < \text{LHS (1.1)} ,$$

a contradiction to the fact that the adjoint pair  $(I, M)$  satisfies (1.1). Hence  $M$  satisfies (CCL).

- (ii)  $\implies$  (i): Now, let  $M$  satisfy (CCL). Consider any arbitrary  $x, y \in [0, 1]$ . Then either  $n(x) > y$  or  $n(x) \leq y$ .

If  $n(x) > y$ , then by Lemma 2.5 (ii) we see that  $M(x, y) = 0$  and hence

$$\text{LHS (1.1)} = I(x, M(x, y)) = I(x, 0) = n(x) = \max(n(x), y) = \text{RHS (1.1)} .$$

If  $n(x) \leq y$  and  $M(x, y) = 0$  then by Lemma 3.1(i) we have that  $n(x) \geq y$  and hence  $n(x) = y$  and it reduces to the above case. Hence let  $M(x, y) > 0$ . Then

$$\text{RHS (1.1)} = \max(n(x), y) = y .$$

We claim now that  $\text{LHS (1.1)} = I(x, M(x, y)) = y$ . If this were not true, then there exists  $1 \geq z > y$  ( $z \not\leq y$  by the monotonicity of  $M$ ) such that

$$I(x, M(x, y)) = \sup\{t \in [0, 1] \mid M(x, t) \leq M(x, y)\} = z .$$

This implies that there exists a  $w \in (0, 1)$  such that  $z > w > y$  and  $M(x, w) \leq M(x, y)$ , which by the monotonicity of t-subnorm implies

that  $M(x, w) = M(x, y)$  with  $w \succ y$ , a contradiction to the fact that  $M$  satisfies (CCL). Thus the adjoint pair  $(I, M)$  satisfies (1.1). □

**Example.** Consider the product t-norm  $T_{\mathbf{P}}(x, y) = xy$ , which is a strict t-norm and hence continuous and Archimedean, whose residual is the Goguen implication given by

$$I_{\mathbf{GG}}(x, y) = \begin{cases} 1, & \text{if } x \leq y, \\ \frac{y}{x}, & \text{if } x > y. \end{cases}$$

It can be easily verified that the pair  $(T_{\mathbf{P}}, I_{\mathbf{GG}})$  does indeed satisfy (1.1) whereas the natural negation of  $T_{\mathbf{P}}$  is the Gödel negation

$$n_{T_{\mathbf{P}}}(x) = I_{\mathbf{GG}}(x, 0) = \begin{cases} 1, & \text{if } x = 0, \\ 0, & \text{if } x > 0. \end{cases}$$

This example clearly shows that the remark of U.Höhle, Remark 1.2, is not always true. In the following we give an equivalence condition under which it is true.

**Theorem 3.2.** Let  $T$  be a continuous t-norm that satisfies (1.1) along with its residual. Then the following are equivalent:

- (i)  $T$  is nilpotent.
- (ii)  $n_T$  is strong.

*Proof.* (i)  $\implies$  (ii): If  $T$  is nilpotent then it is isomorphic to the Łukasiewicz t-norm, i.e., there exists an increasing bijection  $\varphi : [0, 1] \rightarrow [0, 1]$  such that  $T(x, y) = \varphi^{-1}(\max(\varphi(x) + \varphi(y) - 1, 0))$ . It can be easily verified that  $n_T(x) = \varphi^{-1}(1 - \varphi(x))$  which is an involutive negation.

- (ii)  $\implies$  (i): If  $T$  is continuous and satisfies (1.1) along with its residual then, from Theorem 3.1,  $T$  is conditionally cancellative and hence necessarily Archimedean by [9], Proposition 2.15 (ii). Thus  $T$  is either nilpotent or strict. If  $T$  is continuous with a strong natural negation, clearly,  $T$  has zero-divisors and hence  $T$  is nilpotent. □

#### 4. Conditional Cancellativity and Unit element

From the above remarks we note that when the natural negation of the underlying conjunction (a continuous t-norm, in the above case) is strong the class of conjunctions that satisfy (1.1) along with its residual gets restricted. Hence we study the class of t-subnorms  $M$  that satisfy (1.1) along with its residual and whose natural negations are strong. In other words, we seek the characterization of the class of conditionally cancellative t-subnorms with strong natural negations.

Let us recall from the remark following Definition 2.4 that the natural negation of a t-subnorm  $n_M$  need not be a fuzzy negation. If a t-subnorm has 1 as its neutral element, i.e., if it is a t-norm, then we have

$$M(1, y) = 0 \iff y = 0,$$

$$\text{i.e., } y = \sup\{t \mid M(1, t) = 0\} = n_M(1) = 0.$$

Equivalently, by the monotonicity of  $M$  we have that  $n_M$  is a fuzzy negation. However, this is only a necessary and not a sufficient condition.

It was Jenei [6] who proposed some sufficiency conditions and showed that left-continuous t-subnorms with strong natural negations are t-norms, i.e., 1 does become a neutral element.

In the following we show that if a conditionally cancellative t-subnorm is such that  $M(1, y) = y$  for *some*  $y \in (0, 1]$  and if the associated negation is a fuzzy negation then 1 is a neutral element of  $M$ , i.e.,  $M$  is a t-norm. Based on this, we show that in the case  $n_M$  is a strong negation then  $M$  always is a t-norm. In other words, there does not exist any conditionally cancellative *proper* t-subnorm whose natural negation is involutive.

**Lemma 4.1.** *Let  $M$  be a conditionally cancellative t-subnorm. Let  $M(1, y_0) = y_0$ , for some  $y_0 \in (0, 1]$ .*

- (i) *Then  $M(1, y) = y$  for all  $y \in [y_0, 1]$ .*
- (ii) *Let  $y^* = \sup\{t \mid M(1, t) = 0\} = n_M(1)$ . Then  $M(1, y) = y$  for all  $y \in (y^*, y_0]$ .*

*Proof.* Let  $M(1, y_0) = y_0$ , for some  $y_0 \in (0, 1]$ .

- (i) Let  $y_0 < y \leq 1$ . Clearly,  $y_0 = M(1, y_0) < M(1, y) \leq y$ . If  $M(1, y) = y' < y$ , then by associativity and conditional cancellativity we have

$$\left. \begin{array}{l} M(M(1, y_0), y) = M(y_0, y) \\ M(M(1, y), y_0) = M(y', y_0) \end{array} \right\} \implies M(y_0, y) = M(y_0, y') \implies y = y',$$

i.e.,  $M(1, y) = y$  for all  $y \geq y_0$ .

- (ii) Let  $y^* < y \leq y_0$ . Clearly,  $y_0 = M(1, y_0) > y \geq M(1, y) = y'$ . If  $M(1, y) = y' < y$ , then, once again, by associativity and conditional cancellativity we have

$$\left. \begin{array}{l} M(M(1, y_0), y) = M(y_0, y) \\ M(M(1, y), y_0) = M(y', y_0) \end{array} \right\} \implies M(y_0, y) = M(y_0, y') \implies y = y',$$

i.e.,  $M(1, y) = y$  for all  $y \in (y^*, y_0]$ .

□

Based on the above result, we now have the following equivalence condition for a conditionally cancellative t-subnorm to be a t-norm:

**Theorem 4.2.** *Let  $M$  be any conditionally cancellative t-subnorm. Then the following are equivalent:*

- (i)  *$M$  is a t-norm.*




FIGURE 2. The conditionally cancellative (proper) t-subnorms  $M_{\mathbf{P}}$  and  $M_{\mathbf{D}}$  from Remark 4.3

(ii)  $n_M$  is a negation and  $M(1, y_0) = y_0$ , for some  $y_0 \in (0, 1]$ .

*Proof.* Sufficiency is obvious. Necessity follows from the fact that if  $n_M$  is a negation then  $y^* = 0$  in Lemma 4.1 above. □

*Remark 4.3.* Note that both the conditions in Theorem 4.2(ii) are mutually exclusive.

(i) Consider the conditionally cancellative (proper) t-subnorm (see Figure 2 (a))

$$M_{\mathbf{P}}(x, y) = \frac{xy}{2}$$

whose associated negation is a negation but  $M(1, y) \neq y$  for any  $y \in (0, 1]$ .

(ii) Consider the conditionally cancellative (proper) t-subnorm (see Figure 2 (b))

$$M_{\mathbf{D}}(x, y) = \begin{cases} x, & \text{if } y = 1 \text{ and } x \in (0.5, 1] \\ y, & \text{if } x = 1 \text{ and } y \in (0.5, 1] \\ 0, & \text{otherwise} \end{cases},$$

whose associated negation

$$N_M(x) = \begin{cases} 1, & \text{if } x \in [0, 1) \\ 0.5, & \text{if } x = 1 \end{cases}$$

is not a fuzzy negation, since  $N_M(1) \neq 0$ . Note however that  $M(1, y) = M(y, 1) = y$  for any  $y \in (0.5, 1]$ .

The final result of this work shows that in the case  $n_M$  is a strong negation then  $M$  always is a t-norm.

**Theorem 4.4.** *Let  $M$  be any conditionally cancellative t-subnorm. If  $n_M$  is a strong natural negation then  $M$  is a t-norm.*

*Proof.* Our approach will be to show that  $M(1, 1) = 1$  and then the result follows easily from Theorem 4.2. Note also that since  $n_M$  is a strong negation, we have that  $n_M(x) = 1 \iff x = 0$  and  $n_M(x) = 0 \iff x = 1$ . Equivalently,  $M(1, x) = 0 \iff x = 0$ .

On the contrary, let us assume that  $M(1, y) < y$  for all  $y \in (0, 1]$ . In particular,  $M(1, 1) = z$  such that  $0 < z < 1$ . Since  $n_M$  is strong, there exists a  $z' \in (0, 1)$  such that  $z = n_M(z')$ . We claim that  $z' = 0$  and hence  $z = 1$ .

If not, then there exists  $0 < z'' < z'$  and by the definition of  $n_M$  we have that  $M(z, z'') = 0$ . Also, by our assumption  $0 < M(1, z'') = z^* < z''$ . Now, by associativity and conditional cancellativity we have

$$\left. \begin{aligned} M(M(1, 1), z'') &= M(z, z'') \\ M(M(1, z''), 1) &= M(z^*, 1) \end{aligned} \right\} \implies M(z, z'') = 0 = M(z^*, 1) \\ \implies z^* = 0,$$

a contradiction. Thus  $z = 1$  and hence we have the result.  $\square$

## 5. Concluding Remarks

In this work we have solved a more generalised version of an open problem of U.Höhle and have shown that the solution gives a characterization of all conditionally cancellative t-subnorms. Further, by proving an equivalence condition for a conditionally cancellative t-subnorm to be a t-norm, we have shown that conditionally cancellative t-subnorms with involutive natural negations are t-norms.

## References

- [1] Baczyński M., Jayaram B. (2008) *Fuzzy Implications*. Vol. 231, Studies in Fuzziness and Soft Computing, Springer-Verlag, Heidelberg.
- [2] Budinčević M., M.S. Kurilić (1998) A family of strict and discontinuous triangular norms. *Fuzzy Sets and Systems* 95:381-384.
- [3] Demirli K., B. De Baets (1999) Basic properties of implicators in a residual framework, *Tatra Mount. Math. Publ.* 16:31-46.
- [4] Fodor J., I. J. Rudas. (2007) On continuous triangular norms that are migrative. *Fuzzy Sets and Systems* 158:1692-1697.
- [5] Jayaram B., Mesiar R. (2009) *I-Fuzzy Equivalence Relations and I-Fuzzy Partitions* *Info Sci.* 179:1278-1297.
- [6] Jenei S. (2001) Continuity of left-continuous triangular norms with strong induced negations and their boundary condition. *Fuzzy Sets and Systems* 124:35-41.



- [7] Jenei S. (2006) On the convex combination of left-continuous t-norms *Aequationes Math.* 72:47–59.
- [8] Funda Karaçal. (2005) An answer to an open problem on triangular norms. *Fuzzy Sets and Systems* 155(3):459–463.
- [9] Klement E.P., Mesiar R., Pap E. (2000) *Triangular norms*. Kluwer, Dordrecht.
- [10] Klement E.P., Mesiar R., Pap E. (2004) Problems on triangular norms and related operators. *Fuzzy Sets and Systems* 145:471–479.
- [11] Klement E.P., Mesiar R., Pap E. (2006) Open problems posed at the eighth international conference on fuzzy set theory and applications. *Kybernetika* 42:225–235.
- [12] R. Mesiar and V. Novk (1996) Open problems from the 2nd international conference on fuzzy sets theory and its applications. *Fuzzy Sets and Systems* 81:185-190.
- [13] Mesiarová A. (2006) A note on two open problems of Alsina, Frank and Schweizer *Aequationes Math.* 72:41–46.
- [14] Monserrat M., J. Torrens. (2002) On the reversibility of uninorms and t-operators. *Fuzzy Sets and Systems* 131(3):303–314
- [15] Ouyang Y., J. Li (2005) An answer to an open problem on triangular norms. *Info Sci.* 175:78–84.
- [16] Ouyang Y., J. Li, J. Fang (2006) A conditionally cancellative left-continuous t-norm is not necessarily continuous. *Fuzzy Sets and Systems* 157:2328–2332.
- [17] Ouyang Y., J. Li, J. Fang (2007) Further properties of reversible triangular norms. *Fuzzy Sets and Systems* 158:2504–2509.
- [18] Petrík M., P. Sarkoci. (2009) Convex combinations of nilpotent triangular norms. *Jl. Math. Anal. Appln.* 350(1):271–275.
- [19] Ricci R.G. (2009) Solution of an open problem of Alsina, Frank and Schweizer *Aequationes Math.* 77:237–242.
- [20] Sarkoci P.(2008) Dominance is not transitive on continuous triangular norms. *Aequationes Math.* 75:201–207.
- [21] Saminger-Platz S., B. De Baets, H. De Meyer. (2008) A generalization of the Mulholland inequality for continuous Archimedean t-norms. *Jl. Math. Anal. Appln.* 345(2):607–614.
- [22] Viceník P. (2005) Additive generators of associative functions. *Fuzzy Sets and Systems* 153:137-160.
- [23] Viceník P. (2008) Intersections of ranges of additive generators of associative functions. *Tatra Mt. Math. Publ.* 40:117-131
- [24] Viceník P. (2008) Additive generators of border-continuous triangular norms. *Fuzzy Sets and Systems* 159:1631-1645.

Balasubramaniam Jayaram  
 Department of Mathematics,  
 Indian Institute of Technology Hyderabad,  
 Yeddumailaram - 502 205,  
 INDIA  
 e-mail: jbala@iiith.ac.in